



Research Article

Bias Correction for Forecasting PM_{2.5} Concentrations Using Measurement Data from Monitoring Stations by Region

Young Sung Ghim*, Yongjoo Choi, Soontae Kim¹⁾, Chang Han Bae¹⁾, Jinsoo Park²⁾, Hye Jung Shin²⁾

Department of Environmental Science,
Hankuk University of Foreign Studies,
Yongin, Gyeonggi 17035, Republic of
Korea

¹⁾Department of Environmental and
Safety Engineering, Ajou University,
Suwon, Gyeonggi 16499, Republic of
Korea

²⁾Air Quality Research Division, National
Institute of Environmental Research,
Seo, Incheon 22689, Republic of Korea

*Corresponding author.

Tel: +82-31-330-4993

E-mail: ysghim@hufs.ac.kr

Received: 31 July 2018

Revised: 14 September 2018

Accepted: 21 October 2018

ABSTRACT The model and forecasting performances were evaluated to investigate the effectiveness of bias correction for forecasting PM_{2.5} concentrations for the period May 2012 to December 2014. Measured concentrations of PM_{2.5} and major components were obtained from five monitoring stations by region in the Korean Peninsula, and predicted concentrations were obtained from PM_{2.5} simulations using WRF model v3.4.1 and the CMAQ modeling system v4.7.1. Underestimation was prevalent at all stations for all components except NO₃⁻. The effect of bias correction was pronounced at the Gangwon station, where the difference in PM_{2.5} between measured and predicted concentrations was largest. The performances for SO₄²⁻ and the unresolved other component were primarily improved, whereas the performance for NO₃⁻, which was originally overestimated, was degraded. The accuracy of the four-level forecast was moderate at 58% overall, but the probability of detection (POD) of high-concentration events was low at 23%. Bias correction improved the accuracy and POD to 68% and 52%, respectively; however, the rate of false detection of high-concentration events increased as well.

KEY WORDS CMAQ/WRF, Major components, Mean fractional bias, Ratio adjustment, Forecasting performance

1. INTRODUCTION

Concern about PM_{2.5} (particulate matter with an aerodynamic diameter of 2.5 μm or less) has prevailed over the Korean society in the last few years. High concentrations of 24-h PM_{2.5} exceeding 100 μg/m³ at the beginning of 2013 are presumed to have triggered public attention since they followed record-high 1-h averages approaching 1000 μg/m³ in Beijing (Shimadera *et al.*, 2014; Wang *et al.*, 2014; Zhang *et al.*, 2014). Public worries were intensified when the International Agency for Research on Cancer (IARC, 2013) designated PM, as a representative outdoor air pollution, a Group 1 carcinogen in the same year. To meet the public demand for immediate information on PM, the Korean Government launched PM₁₀ and PM_{2.5} forecasting in February 2014 and January 2015, respectively.

In Korea, three-dimensional numerical air quality models are used for PM forecasting. The air quality model predicts pollutant concentrations using emissions and meteorological data by specifying initial and boundary conditions. It is theo-

retically superior to the statistical model because it is based on physical and chemical understanding of atmospheric processes, whereas the latter is based on measurement data. We can construct a best initial field using all of the available data and can minimize the uncertainty of boundary conditions enough to enlarge the modeling domain for the air quality model. We can also obtain a fairly good set of meteorological data because the data-using system is well established, having a long history. However, forecasting using the air quality model cannot be more accurate than the emission data which should have restrictions in reproducing the real-world emissions. Furthermore, there is a big difference between model results, representing the mean of a grid that is several kilometers in length and width and several meters in height, and measurement data from a site installed in a densely populated area.

To improve the accuracy of the forecasting using the air quality model, the differences between model results and measurement data should be reduced, which could be accomplished by improving the models, by improving the input data such as emission data, and by correcting the model biases from measurement data. The first option is best in principle, but it takes considerable time and efforts. Although the second option is generally sought, it, like the first one, has limitations in reducing the aforementioned fundamental differences between model results and measurement data. The third option

forces the model results closer to the measurement data. In the previous study, we investigated the differences in the model performance for measurement data from the intensive monitoring station in Seoul, and found that the ratio adjustment using mean values of model results and measurement data was the most effective of the three bias correction methods (Ghim *et al.*, 2017).

In this study, we first examined the model performance for measurement data from five monitoring stations by region (Fig. 1) for three years from May 2012 to December 2014. Three stations in Seoul, Daejeon, and Gwangju are intensive monitoring stations and two stations in Ulsan and Chuncheon are comprehensive monitoring stations. Because the monitoring stations were distributed by region, we were able to estimate the regional characteristics of the model performance. Next, we examined the effects of bias correction on the model performance by station, applying the ratio adjustment method. Finally, we investigated whether the ratio adjustment method was also effective in improving the forecasting performance, as in the model performance.

2. METHODS

2.1 Modeling

A three-dimensional air quality forecasting system consisting of Weather Research and Forecast (WRF)




Fig. 1. Modeling domain consisting of two grids with horizontal resolutions of 27 and 9 km. Five PM_{2.5} monitoring stations are shown on the fine grid: Seoul (SL) at Bulgwang in Seoul (126.93°E, 37.61°N), Chungcheong (CC) at Munhwa in Daejeon (127.41°E, 36.32°N), Honam (HN) at Oryong in Gwangju (126.85°E, 35.23°N), Yeongnam (YN) at Sinjeong in Ulsan (129.31°E, 35.53°N), and Gangwon (GW) at Seoksa in Chuncheon (127.75°E, 37.86°N).

model v3.4.1 (Skamarock and Klemp, 2008), Sparse Matrix Operator Kernel Emissions Processor (SMOKE) v2.1 (<http://www.smoke-model.org>), and the Community Multiscale Air Quality (CMAQ) modeling system v4.7.1 (Byun and Schere, 2006) was used for PM_{2.5} simulation. WRF model simulations were initialized with Global Forecasting System (GFS) data sets. The WRF model results were prepared for daily emission processing and air quality simulations using the Meteorology–Chemistry Interface Processor. The Statewide Air Pollution Research Center version 99 (SAPRC99) and the fifth-generation modal aerosol model (AEROS) were used as the chemical mechanism and aerosol module, respectively, for the CMAQ simulation.

For anthropogenic emissions, the Intercontinental Chemical Transport Experiment-Phase B (INTEX-B) inventory for the year 2006 (Li *et al.*, 2014; Zhang *et al.*, 2009) was used for Northeast Asia, and the Clean Air Policy Support System (CAPSS) inventory for the year 2007 was used for Korea (Lee *et al.*, 2011; Kim *et al.*, 2008). Biogenic emissions were obtained using the Model of Emissions of Gases and Aerosols from Nature (MEGAN) version 2.04 (Guenther *et al.*, 2006). Fig. 1 shows the modeling domain consisting of two grids with horizontal resolutions of 27 and 9 km. There were 15 layers vertically on a sigma coordinate up to 50 kPa with the lowest layer thickness of about 32 m. Default profiles provided with CMAQ were used as the boundary conditions for the coarse grid, and the boundary conditions for the fine grid were updated by the model outputs from the coarse grid.

2.2 Measurements

PM_{2.5} samples were collected on a Teflon filter (Zeflur, Pall) using a well impactor ninety-six (WINS) and a sequential sampler (PMS-103, APM) at a flow rate of 16.7 L/min for 24 hours. Concentrations of PM_{2.5} and inorganic ions were measured using an automated filter weighing system (MTL) equipped with a microbalance (UMX2, Mettler Toledo) and ion chromatography (ICS 2000, Dionex), respectively. PM_{2.5} samples were also collected on a quartz filter (Tissuquartz 2500QAT-UP, Pall) to measure concentrations of organic and elemental carbons using an OCEC analyzer (Sunset). Concentrations of PM_{2.5} and its components were available on 460 days (47%) at Seoul (SL), 410 days (42%) at Chungcheong (CC), 456 days (47%) at Honam (HN), 329 days (34%) at Yeongnam (YN), and 288 days (30%) at Gangwon

(GW), out of 975 days during the study period.

2.3 Model Performance Metrics

The model performance was evaluated using mean fractional bias (MFB), correlation coefficient (R), and the slope and interceptor of best-fitted line between predicted and measured values. MFB is defined by

$$\text{MFB} = \frac{2}{N} \sum_{i=1}^{i=N} \frac{p_i - m_i}{p_i + m_i} \quad (1)$$

where p_i and m_i denote predicted and measured values, respectively, and N denotes the number of data (Boylan and Russell, 2006). We adopted the performance goals and criteria suggested by Boylan and Russell (2006), which denote the levels of accuracy that the best model can achieve and that are acceptable for standard model applications, respectively. They were given as:

$$\begin{aligned} |\text{MFB}(\text{goals})| &\leq 1.7 e^{-2\bar{C}} + 0.3 \\ |\text{MFB}(\text{criteria})| &\leq 1.4 e^{-2\bar{C}} + 0.6 \end{aligned} \quad (2)$$

where \bar{C} is $(\bar{p} + \bar{m})/2$ in $\mu\text{g}/\text{m}^3$, and \bar{p} and \bar{m} are the means of predicted and measured values, respectively.

2.4 Forecasting Performance Statistics

During the study period, PM_{2.5} concentrations were forecasted by dividing them into four levels as follows: good ($\leq 15 \mu\text{g}/\text{m}^3$), moderate ($15\text{--}50 \mu\text{g}/\text{m}^3$), bad ($50\text{--}100 \mu\text{g}/\text{m}^3$), serious ($> 100 \mu\text{g}/\text{m}^3$). The forecasting performance was evaluated by examining whether the predicted level agreed with the measured level. We distinguished four groups from “A” to “D” and another four groups from “e” to “h” (Fig. 2). “A” indicates that low measured concentrations, which fall into either the good or moderate level, were predicted as high concentrations, which fall into either the bad or serious level. “B” indicates that high measured concentrations were correctly predicted. “e” to “h” indicates that each level was correctly predicted. Because both predicted and measured concentrations were divided into either high or low concentrations (in “A” to “D”), the sum of “A” to “D” is 100%.

We defined four parameters—the accuracy, probability of detection (POD), false alarm rate (FAR), and bias ratio—as shown in Fig. 2 (NIER, 2014; McKeen *et al.*, 2005; USEPA, 2003). The accuracy is the percent of forecasts that correctly predicted the concentration levels. The remaining three parameters examine the quali-

ty of high-concentration forecasts. POD represents the ability to correctly predict high-concentration events, whereas FAR is the percent of high-concentration predictions that did not occur. The bias ratio is the ratio of predicted high-concentration events to observed high-concentration events. A bias ratio greater than 1 indicates that high-concentration events are overpredicted.

3. RESULTS AND DISCUSSION

3.1 Model Performance

Fig. 3 compares the major components between mea-




Fig. 2. Definition of parameters for forecasting performance statistics.




Fig. 3. The mean values of measured and predicted concentrations of major components at monitoring stations by region. The sum of the components is PM_{2.5}, and the other is the remainder of PM_{2.5} excluding the components shown in the figure.

sured and predicted concentrations by station. The measured PM_{2.5} concentration is lowest at YN and highest at GW. YN exhibits the lowest concentrations of major components except for the unresolved other component whose concentration is highest. At GW, the OC concentration is highest whereas the secondary ions are lowest except for YN. Underestimation of the predicted PM_{2.5} is remarkable at GW (Table 1), where the measured PM_{2.5} is highest. In Table 1, the ratio of predicted to measured concentration for NH₄⁺ is close to one on the whole. However, the ratio for NO₃⁻ is greater than 1, and the ratios for carbonaceous components (OC and EC) are less than 0.3, indicating significant underestimation. For the major ions, the overestimation of NO₃⁻ at YN and the underestimation of SO₄²⁻ at GW are notable. At GW, OC underestimation is serious, and EC is underestimated similarly to CC and HN. Despite a substantial overestimation of NO₃⁻ at YN, its effect on PM_{2.5} is insignificant because of a low proportion (Fig. 3). Given a prevalence of underestimations, the model performance is better at YN and SL because of high ratios of predicted to measured concentration for major components.

Table 1. The ratios of predicted to measured concentration for major components by station.

	PM _{2.5}	SO ₄ ²⁻	NO ₃ ⁻	NH ₄ ⁺	OC	EC	Other
SL	0.72	0.59	1.31	0.96	0.44	0.42	0.63
CC	0.66	0.54	1.42	0.93	0.27	0.18	0.57
HN	0.61	0.50	1.45	0.84	0.24	0.17	0.58
YN	0.76	0.55	2.38	1.08	0.26	0.30	0.78
GW	0.52	0.47	1.62	0.87	0.22	0.19	0.33
Overall	0.65	0.53	1.64	0.93	0.29	0.25	0.58

3.2 Bias Correction

Table 2 shows the differences in model performance metrics by station due to bias correction. The predicted concentrations become identical to the measured concentrations because the bias was corrected by multiplying the predicted concentration by the ratio of measured to predicted concentration (ratio adjustment). In contrast, R and the relative intercept remain unchanged. On the whole, the model performance was improved by the bias correction, as MFB moves within the goals from outside and the slope increases from 0.62 to 0.95. The effect of bias correction is noticeable at GW, where MFB moves within the goals from outside the criteria and the

slope increases from the lowest at 0.47 to 0.89. At HN, MFB falls within the goals after correction, but its absolute value is still highest along with that at CC, and the slope increases above 1.0, indicating that the correction effect is unclear.

Table 3 shows the differences in MFB for major components. Overall, MFBs for SO_4^{2-} and the other, which fall outside the criteria and goals, respectively, move within the goals. In contrast, MFB for NO_3^- is pushed outside the criteria because of the correction. MFBs for both OC and EC are improved, but are still outside the criteria. Looking into the differences by station, SO_4^{2-} improves at all stations, as does the other at all stations

Table 2. Differences in model performance metrics by station resulting from bias correction using $\text{PM}_{2.5}$ mean values.

	Measured $\text{PM}_{2.5}$ (mg/m ³)	Predicted	MFB ^a	R	Slope	Relative intercept ^b
(a) Original						
SL	27.3	19.8	-0.36*	0.68	0.58	0.20
CC	28.2	18.6	-0.51*	0.69	0.67	-0.01
HN	26.1	16.0	-0.58*	0.81	0.79	-0.29
YN	24.0	18.1	-0.32*	0.59	0.67	0.12
GW	28.6	14.9	-0.67	0.73	0.47	0.11
Overall	26.8	17.6	-0.48*	0.69	0.62	0.05
(b) Bias corrected						
SL	27.3	27.3	-0.06**	0.68	0.80	0.20
CC	28.2	28.2	-0.13**	0.69	1.01	-0.01
HN	26.1	26.1	-0.13**	0.81	1.29	-0.29
YN	24.0	24.0	-0.06**	0.59	0.88	0.12
GW	28.6	28.6	-0.09**	0.73	0.89	0.11
Overall	26.8	26.8	-0.10**	0.69	0.95	0.05

^a** and * indicate within the goals and criteria, respectively.

^bThe intercept divided by the mean of the predicted values.

Table 3. Differences in mean fractional bias for major components by station resulting from bias correction using $\text{PM}_{2.5}$ mean values.

	SO_4^{2-}	NO_3^-	NH_4^+	OC	EC	Other
(a) Original						
SL	-0.54*	0.52*	-0.06**	-0.77	-0.77	-0.31*
CC	-0.59*	0.48*	-0.13**	-1.16	-1.35	-0.37*
HN	-0.63	0.47*	-0.23**	-1.26	-1.36	-0.42*
YN	-0.65	0.77	-0.08**	-1.10	-0.85*	-0.06**
GW	-0.66	0.65	-0.14**	-1.21	-1.17	-0.77
Overall	-0.61	0.56*	-0.13**	-1.09	-1.10	-0.38*
(b) Bias corrected						
SL	-0.26**	0.74	0.23**	-0.50*	-0.49*	-0.04**
CC	-0.23**	0.76	0.25**	-0.88	-1.10	-0.01**
HN	-0.20**	0.80	0.22**	-0.93	-1.06	0.00**
YN	-0.42*	0.94	0.17**	-0.91	-0.65**	0.19**
GW	-0.10**	1.03	0.43*	-0.76	-0.73*	-0.25**
Overall	-0.25**	0.84	0.25**	-0.79	-0.81	-0.02**

** and * indicate within the goals and criteria, respectively.

except for YN where MFB originally fell within the goals. On the other hand, NO₃⁻ and NH₄⁺ exhibit degradation at all stations; particularly, MFBs for NO₃⁻ at SL, CC and HN move outside the criteria, and MFB for NH₄⁺ at GW moves outside the goals.

3.3 Forecasting Performance

Fig. 4 shows a plot of predicted vs. measured concentrations for PM_{2.5} at all stations. Individual values are compared, different from comparing mean values in the previous sections to examine the model performance. Originally, more data points lie below the 1:1 line, indicating the tendency of underestimation of predicted concentrations (Fig. 4(a)). However, the data points move upward due to bias correction, and the amount of data whose predicted level coincides with the measured level increases, despite some overpredicted data points.

The differences in the forecasting performance statistics that resulted from bias correction are summarized in Table 4. Originally, the overall accuracy for all levels was moderate at 58%, but POD for high-concentration events was only 23% (Table 4(a)). FAR and the bias ratio are also low at 33% and 34%, respectively. High-concentration forecasts are generally fewer, particularly at GW, and consequently, high FAR and the bias ratio at

YN are distinguished. Table 4(b) shows the bias corrected performances. Overall, the accuracy and POD increase by 10% and 30%, respectively, whereas FAR also increases to 56%. Most of all, because the frequency of high-concentration forecasts greatly increases, the bias ratios exceed 100% except for SL. By station, all four parameters at HN and GW greatly increase. The differences in performances between the stations are

Table 4. Differences in forecasting performance statistics^a (%) by station resulting from bias correction using PM_{2.5} mean values.

	Accuracy	POD	FAR	Bias ratio
(a) Original				
SL	61	20	41	33
CC	55	24	27	33
HN	56	39	15	46
YN	65	31	64	85
GW	51	10	0	10
Overall	58	23	33	34
(b) Bias corrected				
SL	69	45	54	98
CC	66	44	61	113
HN	69	79	50	157
YN	70	46	73	169
GW	65	54	49	105
Overall	68	52	56	118

^a See Fig. 2 for the definition of the parameters.




Fig. 4. Plot of predicted vs. measured PM_{2.5} concentrations at all stations. Dotted lines denote the division of concentration levels, and solid lines denote the division of high and low concentrations (see Fig. 2 and the description in the text for details). “Correct” and “false” in the legend indicate that the predicted level coincides and does not coincide with the measured level, respectively. The biases were corrected using mean values by station in (b).

generally reduced by the bias correction, although a high value of POD at HN becomes even higher. A representative case is GW where FAR increases from 0% to 49%.

Note that mean values during the same period were used for bias correction, which cannot be accomplished in the real-time forecasting. However, we tested this bias correction because the effectiveness of bias correction using mean values did not depend much on the period of data used for the correction in our previous study (Ghim *et al.*, 2017). It was probably because the biases of model results from measurement data in Korea were systematically caused by limitations in reproducing the atmospheric environment such as meteorology and emissions during model simulation. The present study revealed that the biases were specific to station (or region) and that the correction should be conducted by station (or region).

4. SUMMARY AND CONCLUSIONS

The model performances and forecasting performances were evaluated using mean and individual data, respectively, for PM_{2.5} and major components from five monitoring stations by region for the period May 2012 to December 2014. WRF model v3.4.1 and the CMAQ modeling system v4.7.1 were used for PM_{2.5} simulation. The effects of bias correction on the two performances were investigated in the second step.

MFB at GW fell outside the criteria because of the lowest predicted concentration despite having the highest measured concentration, whereas those at YN and SL were close to the goals. For the major components, MFBs for NH₄⁺ at all stations fell within the goals. On the other hand, MFB for OC at all stations fell outside the criteria, and MFBs for EC and SO₄²⁻ also performed poorly as they fell outside the criteria at many stations.

The effect of bias correction was pronounced at GW, which had the largest absolute MFB and the smallest slope of the best-fit line, but the performance was improved more than the average for the five stations after correction. In contrast, the effect of correction was unclear at HN, considering that the absolute MFB was still the largest with CC, and the slope increased above 1.0. The performances of SO₄²⁻ and the unresolved other component were improved primarily, whereas the performance of NO₃⁻, which was originally overestimated, was degraded.

The accuracy of the four-level forecast was moderate, at 58% overall; both POD and FAR were low at 23% and 33%, respectively. This tendency was particularly severe at GW, with a POD of 9.8% and a FAR of 0%. Overall, bias correction improved the accuracy and POD to 68% and 52%, respectively, but FAR also increased to 56%. In addition, the differences in performances between stations were generally reduced as POD and FAR at GW greatly increased.

ACKNOWLEDGEMENT

This study was supported by the PM_{2.5} Research Center supported by the Ministry of Science, ICT, and Future Planning (MSIP) and the National Research Foundation (NRF) of Korea (NRF-2014M3C8A5030623), the National Strategic Project-Fine Particle of the National Research Foundation of Korea funded by the Ministry of Science and ICT, the Ministry of Environment, and the Ministry of Health and Welfare (2017M3D8A1092015), and the Hankuk University of Foreign Studies Research Fund.

REFERENCES

- Boylan, J.W., Russell, A.G. (2006) PM and light extinction model performance metrics, goals, and criteria for three-dimensional air quality models. *Atmospheric Environment* 40, 4946–4959.
- Byun, D.W., Schere, K.L. (2006) Review of the governing equations, computational algorithms, and other components of the Models-3 Community Multiscale Air Quality (CMAQ) modeling system. *Applied Mechanics Reviews* 59, 51–77.
- Ghim, Y.S., Choi, Y., Kim, S., Bae, C.H., Park, J., Shin, H.J. (2017) Model performance evaluation and bias correction effect analysis for forecasting PM_{2.5} concentrations. *Journal of Korean Society for Atmospheric Environment* 33, 11–18. (in Korean with English abstract)
- Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P.I., Geron, C. (2006) Estimates of global terrestrial isoprene emissions using MEGAN (Model of emissions of gases and aerosols from nature). *Atmospheric Chemistry and Physics* 6, 3181–3210.
- IARC (International Agency for Research on Cancer) (2013) Outdoor air pollution a leading environmental cause of cancer deaths. Press Release No. 221.
- Kim, S., Moon, N., Byun, D.W. (2008) Korea emissions inventory processing using the US EPA's SMOKE System. *Asian Journal of Atmospheric Environment* 2, 34–46.

- Lee, D.G., Lee, Y.M., Jang, K.W., Yoo, C., Kang, K.H., Lee, J.H., Jung, S.W., Park, J.M., Kee, S.B., Han, J.S., Hong, J.H., Lee, S.J. (2011) Korean national emissions inventory system and 2007 Air pollutant emissions. *Asian Journal of Atmospheric Environment* 5, 278–291.
- Li, M., Zhang, Q., Streets, D.G., He, K.B., Cheng, Y.F., Emmons, L.K., Huo, H., Kang, S.C., Lu, Z., Shao, M., Su, H., Yu, X., Zhang, Y. (2014) Mapping Asian anthropogenic emissions of non-methane volatile organic compounds to multiple chemical mechanisms. *Atmospheric Chemistry and Physics* 14, 5617–5638.
- McKeen, S., Wilczak, J., Grell, G., Djalalova, I., Peckham, S., Hsie, E.Y., Gong, W., Bouchet, V., Moffet, R., McHenry, J., McQueen, J., Tang, Y., Carmichael, G.R., Pagowski, M., Chan, A., Dye, T., Frost, G., Lee, P., Mathur, R. (2005) Assessment of an ensemble of seven real-time ozone forecasts over eastern North America during the summer of 2004. *Journal of Geophysical Research* 110, D21307.
- NIER (National Institute of Environmental Research) (2014) Study on Optimization of the Forecasting Model for Particulate Matter. Prepared by Inha University, Enitech, and Yeungnam University (in Korean).
- Shimadera, H., Hayami, H., Ohara, T., Morino, Y., Takami, A., Irie, S. (2014) Numerical simulation of extreme air pollution by fine particulate matter in China in Winter 2013. *Asian Journal of Atmospheric Environment* 8, 25–34.
- Skamarock, W.C., Klemp, J.B. (2008) A time-split nonhydrostatic atmospheric model for weather research and forecasting applications. *Journal of Computational Physics* 227, 3465–3485.
- USEPA (United States Environmental Protection Agency) (2003) Guidelines for Developing an Air Quality (Ozone and PM_{2.5}) Forecasting Program. Research Triangle Park, NC.
- Wang, Z.F., Li, J., Wang, Z., Yang, W.Y., Tang, X., Ge, B.Z., Yan, P.Z., Zhu, L.L., Chen, X.S., Chen, H.S., Wang, W., Li, J.J., Liu, B., Wang, X.Y., Wang, W., Zhao, Y.L., Lu, N., Su, D.B. (2014) Modeling study of regional severe hazes over Mid-eastern China in January 2013 and its implications on pollution prevention and control. *Science China Earth Sciences* 57, 3–13.
- Zhang, Q., Streets, D.G., Carmichael, G.R., He, K.B., Huo, H., Kannari, A., Klimont, Z., Park, I.S., Reddy, S., Fu, J.S., Chen, D., Duan, L., Lei, Y., Wang, L.T., Yao Z.L. (2009) Asian emissions in 2006 for the NASA INTEX-B mission. *Atmospheric Chemistry and Physics* 9, 5131–5153.
- Zhang, J.K., Sun, Y., Liu, Z.R., Ji, D.S., Hu, B., Liu, Q., Wang, Y.S. (2014) Characterization of submicron aerosols during a month of serious pollution in Beijing, 2013. *Atmospheric Chemistry and Physics* 14, 2887–2903.